

**ARIZONA GAME AND FISH DEPARTMENT  
HERITAGE DATA MANAGEMENT SYSTEM**

**Animal Abstract**

**Element Code:** AFCJC02100

**Data Sensitivity:** No

**CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE**

**NAME:** *Catostomus insignis*

**COMMON NAME:** Sonora Sucker, Gila Sucker

**SYNONYMS:** *Minomus insignis*, *Catostomus insigne*, *Catostomus gila*

**FAMILY:** Catostomidae

**AUTHOR, PLACE OF PUBLICATION:** Baird and Girard, 1854. Descriptions of new species of fishes collected in Texas, New Mexico, and Sonora, by Mr. John H. Clark on the U.S. and Mexican Boundary Survey, and in Texas by Capt. Stewart Van Vliet, U.S.A., Proc. Soc. Nat. Sci. Philadelphia. 7:28.

**TYPE LOCALITY:** Baird and Girard (1854): "Rio San Pedro of the Rio Gila" (San Pedro River, probably near mouth of Babacomari River, Cochise County, Arizona).

**TYPE SPECIMEN:**

**TAXONOMIC UNIQUENESS:** There are 16 other species of the genus in North America.

**DESCRIPTION:** A medium-sized catostomid fish, although adults can attain a size of 80.0 cm (31.5 in). Minckley (1973) states that adults can weigh greater than 2.0 kg (4.4 lbs). "Body fusiform, chubby. Head large. Lower lips enlarged, but only moderately, fleshy lobes not produced. Dorsal fin generally square on distal margin, usually with 11 (rarely 12) fin-rays. Scales relatively large, typically fewer than 60 in lateral line, crowded anteriorly, but not markedly so.

Body sharply bi-colored, brownish dorsally, yellow beneath. Dorso-lateral scales sharply outlined with melanophores over-all, each scale with a discrete broadening of the outline, to form a variably distinct spot; spots aligned to provide a visual effect of longitudinal, punctuate lines on upper sides of darkly-colored individuals. Interradials of fins variably darkened; lower fins typically yellow to white" (Minckley 1973).

"Tiny young of this species, and of most other suckers, have dorsal mouths that migrate to the ventral position as the fish develops through their larval stages" (Stewart 1926 in Minckley 1973).

**AIDS TO IDENTIFICATION:** Very similar in appearance to the Yaqui Sucker, *C. bernardini*. Hybrids have been reported by Clarkson and Minckley (1988) between *C. insignis*

and *C. (Pantosteus) clarki*. Melanophoric spots formed on upper body scales form apparent dash lines. Sometimes this sucker is sharply bicolored.

**ILLUSTRATIONS:**

B&W photo (Minckley 1973:160)  
Line drawing (Page and Burr 1991:170)  
Color drawing (Page and Burr 1991:171)  
Color photos (Rinne and Minckley 1991:20)  
Line drawings (Sublette et al. 1990:200)  
B&W photos (Sublette et al. 1990:200)

**TOTAL RANGE:** Gila and Bill Williams systems (Colorado River drainage) New Mexico and Arizona, also in northern Sonora, Mexico. Sublette, et al. (1990) describe the Sonora Sucker as "native to the Gila and San Francisco drainages (except in extreme headwaters)" in New Mexico.

**RANGE WITHIN ARIZONA:** "...widespread in the Gila and Bill Williams river basins in Arizona" (Sublette et al. 1990). Per 1995 AGFD Native Fish Diversity Review, this fish is thought to be rare to absent in the Salt River Canyon, mainly due to predation by flathead catfish (*Pylodictis olivaris*).

**SPECIES BIOLOGY AND POPULATION TRENDS**

**BIOLOGY:** "*C. insignis* seems intolerant of lake conditions" (Minckley 1973), although a few specimens have been collected at Roosevelt Lake, Arizona, during netting and electrofishing surveys of the late 1980s by AGFD. Sublette et al. 1990 describe the Sonora Suckers of Arizona and New Mexico's San Francisco and Gila Rivers as "very sedentary" despite seasonal changes and major flood events.

**REPRODUCTION:** "Spawning begins in late winter and continues through midsummer. The female is usually attended by two males. Eggs are deposited in riffles, fall into the interstices between gravels, and incubate" (Reughard 1920 in Sublette et al. 1990). "They tend to move to smaller streams or onto riffles in larger streams, but a few populations are known to spawn in lakes" (Minckley 1973). Spawning does not appear to be correlated with any specific pattern of stream flow or temperature.

**FOOD HABITS:** "The young feed along the margins of streams, sometimes by the millions, upon tiny crustaceans, protozoans, and other animal and plant groups" (Minckley 1973). Adults are likewise omnivorous, "feeding in early morning and late evening on the aufwuchs assemblage (diatoms and algae) of shallow pools. A significant component of the diet is macroinvertebrates, particularly Ephemeroptera (Clarkson and Minckley 1988), with some coarse sand occasionally ingested" (Sublette et al. 1990).

**HABITAT:** The Sonora sucker is found in a variety of habitats from warm water rivers to trout streams. "It has an affinity for gravelly or rocky pools, or at least for relatively deep, quiet waters" (Minckley 1973). Adults tend to remain near cover in daylight, but move to runs and deeper riffles at night. Young live and utilize runs and quit eddies.

**ELEVATION:** 369 to 2663 m (1,210 to 8,730 ft.) (AGFD, unpublished data accessed 2001).

**PLANT COMMUNITY:**

**POPULATION TRENDS:** "The status of the species is stable in the San Francisco and Gila River drainages, New Mexico" (Sublette et al. 1990). Thought to be lost from the entire Santa Cruz watershed (D. Foster 2005).

**SPECIES PROTECTION AND CONSERVATION**

<b>ENDANGERED SPECIES ACT STATUS:</b>	SC (USDI, FWS 1996) [C2 USDI, FWS 1994]
<b>STATE STATUS:</b>	1B (AGFD SWAP 21012)
<b>OTHER STATUS:</b>	Forest Service Sensitive, USDA, FS Region 3, 2013 Bureau of Land Management Sensitive (USDI, BLM AZ 2000, 2005, 2008, 2010) P, Listed Endangered (Secretaría de Medio Ambiente 2000, 2010) [Listed Endangered, Secretaría de Desarrollo Social 1994]

**MANAGEMENT FACTORS:** Alteration of historic flow regimes and construction of reservoirs have diminished available habitat for Sonoran Sucker. General watershed erosion causing excessive sand deposition in streams has eliminated much pool habitat required by the species. A winter snagging season for anglers currently (1994) exists for this sucker and the desert sucker below Stewart Mountain Dam on the Lower Salt River. This management action was taken as a measure to encourage harvest of the species, as many die during the extremely slow winter water flows.

**PROTECTIVE MEASURES TAKEN:** Two Arizona Game and Fish Department studies are underway which will document current population dynamics of the Sonora sucker in reaches of two central Arizona rivers: "Effects of Fish Snagging on the Lower Salt River" and "Roundtail Chub Study on the Lower Salt/Verde Rivers". Both of these Enhancement Fund studies will be conducted in FY 94-95.

**SUGGESTED PROJECTS:** Distribution and population trend studies within known range. Variability in reproductive success. Suggest that AZGFD actively search for species in Mexico via cooperators (D. Foster 2005).

**LAND MANAGEMENT/OWNERSHIP:** BIA - Fort Apache, Fort McDowell, Salt River Pima, and San Carlos Reservations; BLM - Kingman, Safford, and Tucson Field Offices; BOR - Phoenix Area; NPS - Montezuma Castle National Monument; USFS - Apache-Sitgreaves, Coconino, Coronado, Prescott, and Tonto National Forests; State Land Department; AGFD Black River Lands; Dead Horse Ranch and Red Rock State Parks; TNC - Canelo Hills Cienega, Patagonia-Sonoita Creek, Aravaipa Canyon and Muleshoe Ranch Preserves; Private.

## **SOURCES OF FURTHER INFORMATION**

### **REFERENCES:**

- Arizona Game and Fish Department Native Fish Diversity Review. 1995. Tempe, Arizona.
- Arizona Game and Fish Department. 2012. Arizona's State Wildlife Action Plan 2012-2022. Phoenix, AZ.
- Clarkson R.W. and W.L. Minckley 1988. Morphology and foods of Arizona catostomid fishes: *Catostomus insignis*, *Pantosteus clarki*, and their putative hybrids. *Copeia* 1988: 422-433.
- Foster, D. 2005. Personal communication.
- Minckley, W.L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix. pp. 160-162.
- Page, L.M. and B.M. Burr. 1991. A field guide to freshwater fishes: North America, north of Mexico. Houghton Mifflin Co., Boston. pp. 169-171.
- Rinne, J.N. and W.L. Minckley. 1991. Native fishes of arid lands: a dwindling resource of the desert southwest. U.S. Department of Agriculture, Forest Service, General Technical Report RM-206. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado. pp. 19-20.
- Secretaría de Desarrollo Social. 1994. Diario Oficial de la Federación. p.50.
- Secretaría de Medio Ambiente. 2000. Diario Oficial de la Federación, PROY-NOM-059-ECOL-2000. p. 44.
- Secretaría de Medio Ambiente y Recursos Naturales. 2010. NORMA Oficial Mexicana NOM-059-SEMARNAT-2010, Protección ambiental-Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo.
- Sublette, J.E., M.D. Hatch and M. Sublette. 1990. The fishes of New Mexico. University of New Mexico Press, Albuquerque. pp. 200-202.
- USDA, Forest Service Region 3. 2013. Regional Forester's Sensitive Species List.
- USDI, Bureau of Land Management. 2000. Arizona BLM Sensitive Species List. Instruction Memorandum No. AZ-2000-018.
- USDI, Bureau of Land Management. 2005. Arizona BLM Sensitive Species List.

USDI, Bureau of Land Management Region 2. 2008. Arizona BLM Sensitive Species List.  
USDI, Bureau of Land Management Region 2. 2010. Arizona BLM Sensitive Species List.  
USDI, Fish and Wildlife Service. 1994. Endangered and Threatened Wildlife and Plants;  
Animal Candidate Review for Listing as Endangered or Threatened Species; Proposed  
Rule. Federal Register 59(219):58996.  
USDI, Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants:  
Review of Plant and Animal Taxa that are Candidates for Listing as Endangered or  
Threatened Species. Federal Register 61(40):7596-7613.

**MAJOR KNOWLEDGEABLE INDIVIDUALS:**

Robert W. Clarkson - U.S. Bureau of Reclamation, AZ. E-mail: rclarkson@lc.usbr.gov.  
John N. Rinne - USFS Rocky Mountain Forest and Range Experiment Station, Flagstaff,  
Arizona.

**ADDITIONAL INFORMATION:**

**Revised:** 1994-07-01 (JJW)  
1995-01-29 (KLY)  
1997-03-04 (SMS)  
2001-10-10 (SMS)  
2002-12-04 (RHB)

To the user of this abstract: you may use this entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp